

1999-2000 NCABA Official Ranking

	League Pins/Gms	Tournament Pins/Gms	Total Pins/Gms	COMPOSITE AVERAGE
1. Russell W. Rose	73704/327	1834/9	75538/336	224.815
2. Mark R. Glover	115837/516	2082/9	117919/525	224.607
3. James M. Wolfe	70581/315	2053/9	72634/324	224.179
4. Chris M. Johnson	69109/312	2002/9	71111/321	221.529
5. James C. Arbogast	88621/402	3813/18	92434/420	220.080
6. Michael E. Burchfield	65209/300	1798/9	67007/309	216.851
7. David M. Lumley	63351/294	2368/12	65719/306	214.767
8. Kenneth E. Lowe	106635/496	1818/9	108453/505	214.758
9. William J. Marshall	63682/297	3184/15	66866/312	214.314
10. Gary Parsons	97949/459	1795/9	99744/468	213.128

On August 9, 1998, the NCABA board of directors voted to reinstitute a system that officially recognizes its high average bowlers according to composite average. Such a system, which began in the 1966-1967 season, was last in effect in 1988-1989.

The above ranking was calculated using the following criteria:

- The bowler must participate in at least three NCABA-sanctioned leagues. For the participation in the league to be acceptable, the bowler must have a minimum of 66 games.
 - The bowler must have a qualifying average as defined above in at least two different bowling centers.
 - The bowler must have a total of at least 200 games in qualifying leagues.
 - The bowler must participate in all events (team, singles, and doubles) of the current bowling season's NCABA Championship Tournament. All scores bowled in this tournament will be used in calculating the composite average.
 - Composite averages will be calculated by totaling all pins in qualifying leagues and the NCABA Championship Tournament and dividing the total by the sum of the games.
-